
JMIRx Med

Overlay journal for preprints with post-review manuscript marketplace
Volume 3 (2022), Issue 4 ISSN 2563-6316 Editor in Chief: Edward Meinert, MA (Oxon), MSc, MBA,
MPA, PhD, CEng, FBCS, EUR ING

Contents

Peer-Review Reports

Peer Review of "Exercise-Induced Hypoalgesia Following Proprioceptive Neuromuscular Facilitation and Resistance Training Among Individuals With Shoulder Myofascial Pain: Randomized Controlled Trial" (e45034) Jonathan Greenberg.....	2
Peer Review of "Exercise-Induced Hypoalgesia Following Proprioceptive Neuromuscular Facilitation and Resistance Training Among Individuals With Shoulder Myofascial Pain: Randomized Controlled Trial" (e45036) Pattanasin Areedomwong.....	4
Peer Review of "Exercise-Induced Hypoalgesia Following Proprioceptive Neuromuscular Facilitation and Resistance Training Among Individuals With Shoulder Myofascial Pain: Randomized Controlled Trial" (e45060) Anonymous.....	6

Authors' Response to Peer Reviews

Authors' Responses to Peer Reviews of "Exercise-Induced Hypoalgesia Following Proprioceptive Neuromuscular Facilitation and Resistance Training Among Individuals With Shoulder Myofascial Pain: Randomized Controlled Trial" (e45038) Zi-Han Xu, Nan An, Zi-Ru Wang.....	8
--	---

Original Paper

Exercise-Induced Hypoalgesia Following Proprioceptive Neuromuscular Facilitation and Resistance Training Among Individuals With Shoulder Myofascial Pain: Randomized Controlled Trial (e40747) Zi-Han Xu, Nan An, Zi-Ru Wang.....	13
--	----

Peer-Review Report

Peer Review of "Exercise-Induced Hypoalgesia Following Proprioceptive Neuromuscular Facilitation and Resistance Training Among Individuals With Shoulder Myofascial Pain: Randomized Controlled Trial"

Jonathan Greenberg^{1,2}, PhD

¹Department of Psychiatry, Harvard Medical School, Boston, MA, United States

²Massachusetts General Hospital, Boston, MA, United States

Related Articles:

Companion article: <https://www.medrxiv.org/content/10.1101/2022.06.28.22276990v1>

Companion article: <https://preprints.jmir.org/preprint/40747>

Companion article: <https://med.jmirx.org/2022/4/e45038/>

Companion article: <https://med.jmirx.org/2022/4/e40747/>

(*JMIRx Med* 2022;3(4):e45034) doi:[10.2196/45034](https://doi.org/10.2196/45034)

KEYWORDS

exercise induced hypoalgesia; proprioceptive neuromuscular facilitation; PNF; resistance exercise; conditioned pain modulation; myofascial pain syndrome; resistance training; hypoalgesia; exercise-induced hypoalgesia; shoulder myofascial pain; myofascial pain; pain management; chronic pain; musculoskeletal pain; physical therapy; physiotherapy; shoulder pain; upper back pain; exercise; pain

This is a peer-review report submitted for the paper "Exercise-Induced Hypoalgesia Following Proprioceptive Neuromuscular Facilitation and Resistance Training Among Individuals With Shoulder Myofascial Pain: Randomized Controlled Trial"

Round 1 Review

This paper [1] set to estimate the effect of proprioceptive neuromuscular facilitation (PNF) and resistance training on exercise-induced analgesia and conditioned pain modulation among young adult women with myofascial pain syndrome.

The paper holds several strengths, including random assignment and the inclusion of PNF, 2 resistance training exercise types, and 1 passive control group, which enables comparison across exercise conditions. Authors justifiably correct for multiple comparisons. The discussion thoroughly interprets the findings and relates them to existing literature. Some questions and potential limitations are listed below.

Specific Comments

Major Comments

1. The study is limited to young women (18-30 years old) and therefore has limited generalizability to men, as well as women

above the age of 30 years. Authors partially acknowledge this in the limitations section (with regard to gender).

2. The sample size in each group is modest (n=18-20), limiting statistical power.

3. Did the authors have a specific hypothesis about the relative effect of PNF, isometric, and isotonic exercise training on outcomes? Such a hypothesis is now stated. Was the testing of differential effects exploratory?

4. Authors indicate that "Randomized sequences were generated by computer." Can authors provide details on the method, software, or website used for randomization?

5. Authors indicate that participants were excluded if they experienced depression, psychosis, cognitive impairment, etc. How were these assessed?

Minor Comments

6. Authors make use of 6 different acronyms in the abstract, which may make it more difficult to read, particularly for individuals outside this immediate field. When possible, consider spelling things out to increase ease of readability.

7. Please change all instances of " $P<.000$ " to " $P<.001$ "

8. There are some typos throughout the manuscript, please correct those (eg, “Our findings mostly met what we previously hypothesized, where was an increase in PPT at trigger point”; which may have been “There was an increase,” or “Crombie et al investigated that the serum endocannabinoids increased,”

which may have been “Crombie et al ‘found’ or ‘reported’ that....,” as well as other examples throughout).

Round 2 Review

Thank you for revising the manuscript. The authors have addressed my concerns.

Conflicts of Interest

None declared.

Reference

1. Xu ZH, An N, Wang ZR. Exercise-Induced Hypoalgesia Following Proprioceptive Neuromuscular Facilitation and Resistance Training Among Individuals With Shoulder Myofascial Pain: Randomized Controlled Trial. *JMIRx Med* 2022;3(4):e40747 [[FREE Full text](#)]

Abbreviations

PNF: proprioceptive neuromuscular facilitation

Edited by E Meinert; submitted 13.12.22; this is a non-peer-reviewed article; accepted 13.12.22; published 27.12.22.

Please cite as:

Greenberg J

Peer Review of "Exercise-Induced Hypoalgesia Following Proprioceptive Neuromuscular Facilitation and Resistance Training Among Individuals With Shoulder Myofascial Pain: Randomized Controlled Trial"

JMIRx Med 2022;3(4):e45034

URL: <https://med.jmirx.org/2022/4/e45034>

doi: [10.2196/45034](https://doi.org/10.2196/45034)

PMID:

©Jonathan Greenberg. Originally published in *JMIRx Med* (<https://med.jmirx.org>), 27.12.2022. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in the *JMIRx Med*, is properly cited. The complete bibliographic information, a link to the original publication on <https://med.jmirx.org/>, as well as this copyright and license information must be included.

Peer-Review Report

Peer Review of “Exercise-Induced Hypoalgesia Following Proprioceptive Neuromuscular Facilitation and Resistance Training Among Individuals With Shoulder Myofascial Pain: Randomized Controlled Trial”

Pattanasin Areedomwong¹, PhD

Department of Physical Therapy, Mae Fah Luang University, Chiang Rai, Thailand

Related Articles:

Companion article: <https://www.medrxiv.org/content/10.1101/2022.06.28.22276990v1>

Companion article: <https://preprints.jmir.org/preprint/40747>

Companion article: <https://med.jmirx.org/2022/4/e45038/>

Companion article: <https://med.jmirx.org/2022/4/e40747/>

(*JMIRx Med* 2022;3(4):e45036) doi:[10.2196/45036](https://doi.org/10.2196/45036)

KEYWORDS

exercise induced hypoalgesia; proprioceptive neuromuscular facilitation; PNF; resistance exercise; conditioned pain modulation; myofascial pain syndrome; resistance training; hypoalgesia; exercise-induced hypoalgesia; shoulder myofascial pain; myofascial pain; pain management; chronic pain; musculoskeletal pain; physical therapy; physiotherapy; shoulder pain; upper back pain; exercise; pain

This is a peer-review report submitted for the paper “Exercise-Induced Hypoalgesia Following Proprioceptive Neuromuscular Facilitation and Resistance Training Among Individuals With Shoulder Myofascial Pain: Randomized Controlled Trial”

Round 1 Review

General Comments

This paper [1] shows the effects of several types of exercise on exercise-induced hypoalgesia and conditioned pain modulation in patients with myofascial pain syndrome (MPS) of upper trapezius muscle. I thoroughly reviewed this work and find that there is room for improvement. The comments are as follows.

Specific Comments**Major Comments**

1. Why was this sample size chosen?
2. Regarding the first inclusion criteria, the patients must report their MPS for at least 4 weeks up to 3 months. In my opinion, the patients do not know whether they have MPS, and they commonly complain about their shoulder pain only. They will be informed about having MPS after being diagnosed by the physicians. Thus, it is not clear what the statement, “reported MPS” is referring to. In addition, I am not sure if patients with

cardiovascular conditions, such as uncontrolled hypertension, are excluded from the study. Is it because of some risks of cardiovascular problems while performing accidental events during exercises such as holding the breath or due to resistant exercise-induced cardiovascular problems?

3. I am not sure how 60% maximum voluntary contraction (MVC) or pain-free load is set for each exercise intervention. Can we set it with other intensities? Is it possible that the patients doing an exercise with 60% MVC or pain-free load show significant differences on the outcomes? Why does the proprioceptive neuromuscular facilitation (PNF) method use 60% MVC for designing the PNF training protocol? The PNF intensity is set as a maximal resistance of the patients to facilitate an optimal outcome. Some techniques, such as the hold-relax technique with maximal resistance but within subpain threshold, are effectively used for relaxing muscle spasm or guarding in patients with muscle pain conditions. The authors apply agonist reversal, combination of isotonic contraction, and rhythmic stabilization without providing details of start and end positions, thus making it difficult to follow.

Minor Comments**Abstract**

1. Please provide a specific name for a muscle affected by MPS, such as “patients with MPS of upper trapezius muscle.”

2. Exercise-induced hypoalgesia should be defined in relation to what outcomes are included for it.
3. Please specify specific area of pressure pain threshold of remote sites on the extensor carpus radialis and the peroneus longus.
4. Is the visual analog scale one of the study's outcomes?

Conflicts of Interest

None declared.

Reference

1. Xu ZH, An N, Wang ZR. Exercise-Induced Hypoalgesia Following Proprioceptive Neuromuscular Facilitation and Resistance Training Among Individuals With Shoulder Myofascial Pain: Randomized Controlled Trial. *JMIRx Med* 2022;3(4):e40747 [[FREE Full text](#)]

Abbreviations

MPS: myofascial pain syndrome

MVS: maximum voluntary contraction

PNF: proprioceptive neuromuscular facilitation

Edited by E Meinert; submitted 13.12.22; this is a non-peer-reviewed article; accepted 13.12.22; published 27.12.22.

Please cite as:

Areedomwong P

Peer Review of "Exercise-Induced Hypoalgesia Following Proprioceptive Neuromuscular Facilitation and Resistance Training Among Individuals With Shoulder Myofascial Pain: Randomized Controlled Trial"

JMIRx Med 2022;3(4):e45036

URL: <https://med.jmirx.org/2022/4/e45036>

doi: [10.2196/45036](https://doi.org/10.2196/45036)

PMID:

©Pattanasin Areedomwong. Originally published in *JMIRx Med* (<https://med.jmirx.org>), 27.12.2022. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in *JMIRx Med*, is properly cited. The complete bibliographic information, a link to the original publication on <https://med.jmirx.org/>, as well as this copyright and license information must be included.

Peer Review of “Exercise-Induced Hypoalgesia Following Proprioceptive Neuromuscular Facilitation and Resistance Training Among Individuals With Shoulder Myofascial Pain: Randomized Controlled Trial”

Anonymous

Corresponding Author:

Related Articles:

Companion article: <https://www.medrxiv.org/content/10.1101/2022.06.28.22276990v1>

Companion article: <https://preprints.jmir.org/preprint/40747>

Companion article: <https://med.jmirx.org/2022/4/e45038/>

Companion article: <https://med.jmirx.org/2022/4/e40747/>

(*JMIRx Med* 2022;3(4):e45060) doi:[10.2196/45060](https://doi.org/10.2196/45060)

KEYWORDS

exercise induced hypoalgesia; proprioceptive neuromuscular facilitation; PNF; resistance exercise; conditioned pain modulation; myofascial pain syndrome; resistance training; hypoalgesia; exercise-induced hypoalgesia; shoulder myofascial pain; myofascial pain; pain management; chronic pain; musculoskeletal pain; physical therapy; physiotherapy; shoulder pain; upper back pain; exercise; pain

This is a peer-review report submitted for the paper “Exercise-Induced Hypoalgesia Following Proprioceptive Neuromuscular Facilitation and Resistance Training Among Individuals With Shoulder Myofascial Pain: Randomized Controlled Trial”

Round 1 Review

General Comments

This paper [1] aims to compare short-term exercise-induced hypoalgesia responses following different types of exercise in pain modulation within patients with myofascial pain. It is generally well written and presents innovative results to clarify the knowledge in the treatment of myofascial pain.

Specific Comments

Major Comments

1. Methods: In the procedures section, please add information about possible blinding of the evaluators (ie, experience of the persons who did the manual assessment of the myofascial pain syndrome, people who performed the exercise programs, etc).

Minor Comments

2. Discussion: Please try to address the important improvements in the proprioceptive neuromuscular facilitation exercise group in relation to personal interaction with the researcher (manual contact, personal adaptation, etc).

Conflicts of Interest

None declared.

Reference

1. Zu ZH, An N, Wang ZR. Exercise-Induced Hypoalgesia Following Proprioceptive Neuromuscular Facilitation and Resistance Training Among Individuals With Shoulder Myofascial Pain: Randomized Controlled Trial. *JMIRx Med* 2022;3(4):e40747 [[FREE Full text](#)]
-

Edited by E Meinert; submitted 14.12.22; this is a non-peer-reviewed article; accepted 14.12.22; published 27.12.22.

Please cite as:

Anonymous

Peer Review of "Exercise-Induced Hypoalgesia Following Proprioceptive Neuromuscular Facilitation and Resistance Training Among Individuals With Shoulder Myofascial Pain: Randomized Controlled Trial"

JMIRx Med 2022;3(4):e45060

URL: <https://med.jmirx.org/2022/4/e45060>

doi: [10.2196/45060](https://doi.org/10.2196/45060)

PMID:

© Anonymous. Originally published in JMIRx Med (<https://med.jmirx.org>), 27.12.2022. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIRx Med, is properly cited. The complete bibliographic information, a link to the original publication on <https://med.jmirx.org/>, as well as this copyright and license information must be included.

Authors' Response to Peer Reviews

Authors' Responses to Peer Reviews of "Exercise-Induced Hypoalgesia Following Proprioceptive Neuromuscular Facilitation and Resistance Training Among Individuals With Shoulder Myofascial Pain: Randomized Controlled Trial"

Zi-Han Xu¹, MSc; Nan An¹, BSc; Zi-Ru Wang¹, MSc

School of Sport Medicine and Rehabilitation, Beijing Sport University, Beijing, China

Corresponding Author:

Zi-Han Xu, MSc

School of Sport Medicine and Rehabilitation

Beijing Sport University

48 Xinxu Road, Haidian district

Beijing, 100084

China

Phone: 86 17888837585

Email: 2290259311@qq.com

Related Articles:

Companion article: <https://www.medrxiv.org/content/10.1101/2022.06.28.22276990v1>

Companion article: <https://preprints.jmir.org/preprint/40747>

Companion article: <https://med.jmirx.org/2022/4/e45034/>

Companion article: <https://med.jmirx.org/2022/4/e45060/>

Companion article: <https://med.jmirx.org/2022/4/e45036/>

Companion article: <https://med.jmirx.org/2022/4/e40747/>

(*JMIRx Med* 2022;3(4):e45038) doi:[10.2196/45038](https://doi.org/10.2196/45038)

KEYWORDS

exercise induced hypoalgesia; proprioceptive neuromuscular facilitation; PNF; resistance exercise; conditioned pain modulation; myofascial pain syndrome; resistance training; hypoalgesia; exercise-induced hypoalgesia; shoulder myofascial pain; myofascial pain; pain management; chronic pain; musculoskeletal pain; physical therapy; physiotherapy; shoulder pain; upper back pain; exercise; pain

This is the authors' response to peer-review reports for "Exercise-Induced Hypoalgesia Following Proprioceptive Neuromuscular Facilitation and Resistance Training Among Individuals With Shoulder Myofascial Pain: Randomized Controlled Trial"

Round 1 Review

Responses to Comments From Reviewer AC [1]**General Comments**

This paper [2] set to estimate the effect of proprioceptive neuromuscular facilitation (PNF) and resistance training on

exercise-induced analgesia and conditioned pain modulation among young adult women with myofascial pain syndrome.

The paper holds several strengths, including random assignment and the inclusion of PNF, 2 resistance training exercise types, and 1 passive control group, which enables comparison across exercise conditions. Authors justifiably correct for multiple comparisons. The discussion thoroughly interprets the findings and relates them to existing literature. Some questions and potential limitations are listed below.

Response:

We appreciate this supportive feedback.

Major Comments

1. The study is limited to young women (18-30 years old) and therefore has limited generalizability to men, as well as women above the age of 30 years. Authors partially acknowledge this in the limitations section (with regard to gender).

Response:

Thank you for raising this point. We adjusted statements about gender and age in the limitations section. In our university, female students were the most common group of shoulder and neck pain considering the gender difference in pain perception and sedentary behaviors, so we enrolled female students to enhance the comparability of the results.

2. The sample size in each group is modest (n=18-20), limiting statistical power.

Response:

Thanks for your comment. We agree that the sample size is relatively small and may attenuate the statistical power. Since this is a pilot study aiming at investigating the potential relationship, and given the feasibility issues during the national COVID-19 prevention and control policies in our setting, we were trying to recruit as many participants as we could to meet the basic requirement to appeal the effect size. We added more description about the sample size calculation in the participant section.

3. Did the authors have a specific hypothesis about the relative effect of PNF, isometric, and isotonic exercise training on outcomes? Such a hypothesis is now stated. Was the testing of differential effects exploratory?

Response:

Thank you for your valuable and considerate suggestion. Considering the recent progress [3] in the central endogenous pain modulation, we found that the different C afferent fibers' input can trigger the descending inhibition or facilitation pathway. In the context of resistance exercise, the concentric or eccentric muscle contraction with a subpain threshold may activate the nonnoxious C fibers and induce pain inhibition. Thus, we developed and added our preliminary hypothesis in the discussion section and revised the mechanisms of endogenous pain modulation in the introduction section.

4. Authors indicate that "Randomized sequences were generated by computer." Can authors provide details on the method, software, or website used for randomization?

Response:

Thank you for the above suggestions. All participants were labeled from number 01 to 76, then the sequence was randomized using Excel (Microsoft Corp), and allocated following the A-B-C-D circulation order. We added the detailed method of randomization in the participant section.

5. Authors indicate that participants were excluded if they experienced depression, psychosis, cognitive impairment, etc. How were these assessed?

Response:

Thank you for raising this important point. During the screening of participants, our lab employed a physician who was responsible for evaluating the medical risks for all participants, including depression, psychosis, and cognitive impairment. The self-rating depression scale, self-rating anxiety scale, and the brief psychiatric rating scale were also applied and assessed by the physician during the screening periods.

Minor Comments

6. Authors make use of 6 different acronyms in the abstract, which may make it more difficult to read, particularly for individuals outside this immediate field. When possible, consider spelling things out to increase ease of readability.

Response:

We apologize for the lack of clarity. We reduced the acronyms as far as we can in the revised manuscript. Only 3 acronyms including "PNF," "CPM," and "EIH" were retained.

7. Please change all instances of " $P<.000$ " to " $P<.001$ "

Response:

Thank you so much for your careful check; we have changed all reports about the *P* values according to your and our editors' comments.

8. There are some typos throughout the manuscript, please correct those (eg, "Our findings mostly met what we previously hypothesized, where was an increase in PPT at trigger point"; which may have been "There was an increase," or "Crombie et al investigated that the serum endocannabinoids increased," which may have been "Crombie et al 'found' or 'reported' that....," as well as other examples throughout).

Response:

We apologize for the typos. We double-checked and revised all typos and grammar errors in the manuscript.

Responses to Comments From Reviewer Anonymous [4]

General Comments

This paper [2] aims to compare short-term exercise-induced hypoalgesia responses following different types of exercise in pain modulation within patients with myofascial pain. It is generally well written and presents innovative results to clarify the knowledge in the treatment of myofascial pain.

Response:

We thank the reviewer for reading our paper carefully and for the encouraging feedback.

Major Comments

1. Methods: In the procedures section, please add information about possible blinding of the evaluators (ie, experience of the persons who did the manual assessment of the myofascial pain syndrome, people who performed the exercise programs, etc).

Response:

Thank you for raising this point. The allocation of researchers was strictly carried out following the double-blinded principle,

and we separated evaluation and intervention across different researchers. We have added all information about researchers in the procedure section.

Minor Comments

2. Discussion: Please try to address the important improvements in the proprioceptive neuromuscular facilitation exercise group in relation to personal interaction with the researcher (manual contact, personal adaptation, etc).

Response:

Thanks for your valuable comment; we agree that the interaction between the participants and the physical therapist has positive effects on proprioception processing and pain management. We have addressed this mechanism at the end of the “potential mechanism of EIH” part in the Discussion section.

Responses to Comments From Reviewer DA [5]:

General Comments

This paper [2] shows the effects of several types of exercise on exercise-induced hypoalgesia and conditioned pain modulation in patients with myofascial pain syndrome (MPS) of upper trapezius muscle. I thoroughly reviewed this work and find that there is room for improvement. The comments are as follows.

Response:

We thank the reviewer for reading our paper carefully and giving the above positive comments.

Major Comments

1. Why was this sample size chosen?

Response:

Thank you for pointing this out. According to the pressure pain threshold (PPT) changes seen in previous studies, we used G-Power software with *F* test and ANCOVA parameters to calculate the sample size. The total samples of this study should be a minimum of 76 participants in the 3 groups, or 19 participants in each group. We added these sample size calculation methods into the participant section.

2. Regarding the first inclusion criteria, the patients must report their MPS for at least 4 weeks up to 3 months. In my opinion, the patients do not know whether they have MPS, and they commonly complain about their shoulder pain only. They will be informed about having MPS after being diagnosed by the physicians. Thus, it is not clear what the statement, “reported MPS” is referring to. In addition, I am not sure if patients with cardiovascular conditions, such as uncontrolled hypertension, are excluded from the study. Is it because of some risks of cardiovascular problems while performing accidental events during exercises such as holding the breath or due to resistant exercise-induced cardiovascular problems?

Response:

We gratefully thank you for the precious time your spent making constructive remarks.

First, we are very sorry for the inaccurate writing of the inclusion criteria, and it should be “reported shoulder pain” in

practice. During the recruitment period, we brought in all participants reporting shoulder pain, then carefully evaluated their symptoms following the criteria of the upper trapezius MPS. Second, our lab employed a physician who was responsible for evaluating the medical risks for all participants, including cardiovascular problems during the screening periods. Participants who presented hypertension or other risky cardiovascular conditions have been excluded from the study.

3. I am not sure how 60% maximum voluntary contraction (MVC) or pain-free load is set for each exercise intervention. Can we set it with other intensities? Is it possible that the patients doing an exercise with 60% MVC or pain-free load show significant differences on the outcomes? Why does the proprioceptive neuromuscular facilitation (PNF) method use 60% MVC for designing the PNF training protocol? The PNF intensity is set as a maximal resistance of the patients to facilitate an optimal outcome. Some techniques, such as the hold-relax technique with maximal resistance but within subpain threshold, are effectively used for relaxing muscle spasm or guarding in patients with muscle pain conditions. The authors apply agonist reversal, combination of isotonic contraction, and rhythmic stabilization without providing details of start and end positions, thus making it difficult to follow.

Response:

Thank you for your valuable and constructive comments. (1) Considering the previous EIH studies, the optimal intensity of resistance exercise in patients with chronic pain may not be 100% MVC or over-pain thresholds, and it may elicit pain symptoms during the exercise [6]. In most cases, we may prefer exercise with subpain threshold intensity (around 50%-60% MVC) in the context of pain management [7]. (2) For the PNF intervention, it is true that the maximal resistance can facilitate an outcome about flexibility and the joint range of motion. However, some previous studies [8,9] compared the analgesic effects of PNF hold-relax methods with different intensities and found that exercise with 60%-70% MVC may be more suitable for pain treatment. In our study, the intensity of exercise was set as 60% MVC, and then it could adjust to subpain thresholds if participants reported pain during the exercise. (3) In addition, as the reviewer suggested, we have rewritten the PNF method section and added practical details in the agonist reversal, combination of isotonic contraction, and rhythmic stabilization training, including the joint positions and movements.

Minor Comments

1. Please provide a specific name for a muscle affected by MPS, such as “patients with MPS of upper trapezius muscle.”

Response:

Thank you for pointing this out. We specified the upper trapezius as the MPS-affected muscles in the Abstract and Methods sections.

2. Exercise-induced hypoalgesia should be defined in relation to what outcomes are included for it.

Response:

Thank you for the above suggestions. We defined the improvement of PPTs as the EIH response.

3. Please specify specific area of pressure pain threshold of remote sites on the extensor carpus radialis and the peroneus longus.

Response:

Thank you for raising this concern. We have added the specific test point of the extensor carpus radialis (5 cm below the lateral

condyle of humerus) and the peroneus longus (10 cm below the lateral femoral condyle) in the outcome measure section.

4. Is the visual analog scale one of the study's outcomes?

Response:

No. The visual analog scale would be only measured as the baseline characteristics and would not be considered as an outcome measurement. We moved the visual analog scale-related parts to the Procedures section.

References

- Greenberg J. Peer Review of "Exercise-Induced Hypoalgesia Following Proprioceptive Neuromuscular Facilitation and Resistance Training Among Individuals With Shoulder Myofascial Pain: Randomized Controlled Trial". *JMIRx Med* 2022;3(4):e45034 [FREE Full text] [doi: [10.2196/45034](https://doi.org/10.2196/45034)]
- Xu ZH, An N, Wang ZR. Exercise-Induced Hypoalgesia Following Proprioceptive Neuromuscular Facilitation and Resistance Training Among Individuals With Shoulder Myofascial Pain: Randomized Controlled Trial. *JMIRx Med* 2022;3(4):e40747 [FREE Full text] [doi: [10.2196/40747](https://doi.org/10.2196/40747)]
- You HJ, Lei J, Pertovaara A. Thalamus: The 'promoter' of endogenous modulation of pain and potential therapeutic target in pathological pain. *Neurosci Biobehav Rev* 2022 Aug;139:104745. [doi: [10.1016/j.neubiorev.2022.104745](https://doi.org/10.1016/j.neubiorev.2022.104745)] [Medline: [35716873](https://pubmed.ncbi.nlm.nih.gov/35716873/)]
- Anonymous. Peer Review of "Exercise-Induced Hypoalgesia Following Proprioceptive Neuromuscular Facilitation and Resistance Training Among Individuals With Shoulder Myofascial Pain: Randomized Controlled Trial". *JMIRx Med* 2022;3(4):e45060 [FREE Full text] [doi: [10.2196/45060](https://doi.org/10.2196/45060)]
- Areudomwong P. Peer Review of "Exercise-Induced Hypoalgesia Following Proprioceptive Neuromuscular Facilitation and Resistance Training Among Individuals With Shoulder Myofascial Pain: Randomized Controlled Trial". *JMIRx Med* 2022;3(4):e45036 [FREE Full text] [doi: [10.2196/45036](https://doi.org/10.2196/45036)]
- Coombes BK, Wiebusch M, Heales L, Stephenson A, Vicenzino B. Isometric Exercise Above but not Below an Individual's Pain Threshold Influences Pain Perception in People With Lateral Epicondylalgia. *Clin J Pain* 2016 Dec;32(12):1069-1075. [doi: [10.1097/AJP.0000000000000365](https://doi.org/10.1097/AJP.0000000000000365)] [Medline: [26889612](https://pubmed.ncbi.nlm.nih.gov/26889612/)]
- Vaegter HB, Handberg G, Graven-Nielsen T. Similarities between exercise-induced hypoalgesia and conditioned pain modulation in humans. *Pain* 2014 Jan;155(1):158-167. [doi: [10.1016/j.pain.2013.09.023](https://doi.org/10.1016/j.pain.2013.09.023)] [Medline: [24076045](https://pubmed.ncbi.nlm.nih.gov/24076045/)]
- Lim W. Optimal intensity of PNF stretching: maintaining the efficacy of stretching while ensuring its safety. *J Phys Ther Sci* 2018 Aug;30(8):1108-1111 [FREE Full text] [doi: [10.1589/jpts.30.1108](https://doi.org/10.1589/jpts.30.1108)] [Medline: [30154610](https://pubmed.ncbi.nlm.nih.gov/30154610/)]
- Kwak DH, Ryu YU. Applying proprioceptive neuromuscular facilitation stretching: optimal contraction intensity to attain the maximum increase in range of motion in young males. *J Phys Ther Sci* 2015 Jul;27(7):2129-2132 [FREE Full text] [doi: [10.1589/jpts.27.2129](https://doi.org/10.1589/jpts.27.2129)] [Medline: [26310658](https://pubmed.ncbi.nlm.nih.gov/26310658/)]

Abbreviations

EIH: exercise-induced hypoalgesia

MPS: myofascial pain syndrome

MVC: maximum voluntary contraction

PNF: proprioceptive neuromuscular facilitation

PPT: pressure pain threshold

Edited by E Meinert; submitted 13.12.22; this is a non-peer-reviewed article; accepted 13.12.22; published 27.12.22.

Please cite as:

Xu ZH, An N, Wang ZR

Authors' Responses to Peer Reviews of "Exercise-Induced Hypoalgesia Following Proprioceptive Neuromuscular Facilitation and Resistance Training Among Individuals With Shoulder Myofascial Pain: Randomized Controlled Trial"

JMIRx Med 2022;3(4):e45038

URL: <https://med.jmirx.org/2022/4/e45038>

doi: [10.2196/45038](https://doi.org/10.2196/45038)

PMID:

©Zi-Han Xu, Nan An, Zi-Ru Wang. Originally published in JMIRx Med (<https://med.jmirx.org>), 27.12.2022. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIRx Med, is properly cited. The complete bibliographic information, a link to the original publication on <https://med.jmirx.org/>, as well as this copyright and license information must be included.

Original Paper

Exercise-Induced Hypoalgesia Following Proprioceptive Neuromuscular Facilitation and Resistance Training Among Individuals With Shoulder Myofascial Pain: Randomized Controlled Trial

Zi-Han Xu^{1*}, MSc; Nan An^{1*}, BSc; Zi-Ru Wang^{1*}, MSc

School of Sport Medicine and Rehabilitation, Beijing Sport University, Beijing, China

* all authors contributed equally

Corresponding Author:

Zi-Han Xu, MSc

School of Sport Medicine and Rehabilitation

Beijing Sport University

48 Xinxu Road, Haidian district

Beijing, 100084

China

Phone: 86 17888837585

Email: 2290259311@qq.com

Related Articles:

Companion article: <https://www.medrxiv.org/content/10.1101/2022.06.28.22276990v1>

Companion article: <https://preprints.jmir.org/preprint/40747>

Companion article: <https://med.jmirx.org/2022/4/e45034/>

Companion article: <https://med.jmirx.org/2022/4/e45060/>

Companion article: <https://med.jmirx.org/2022/4/e45036/>

Companion article: <https://med.jmirx.org/2022/4/e45038/>

Abstract

Background: Various exercises can attenuate pain perception in healthy individuals and may interact with the descending pain modulation in the central nervous system. However, the analgesic effects of exercise in patients with myofascial pain can be disrupted by the pathological changes during chronic pain conditions. Thus, the exercises targeted on the facilitation of the sensory-motor interaction may have a positive impact on the restoration of the descending pain modulation and the analgesia effects.

Objective: This paper estimates the effect of proprioceptive neuromuscular facilitation (PNF) and resistance training on exercise-induced hypoalgesia (EIH) and conditioned pain modulation (CPM) among patients with myofascial pain syndrome.

Methods: A total of 76 female patients with myofascial pain syndrome (aged 18-30 years), with the pain in the upper trapezius and a visual analog scale score of greater than 30/100 mm, were enrolled in the study. Participants were randomly assigned into 3 intervention groups, including isometric (n=18, 24%), isotonic (n=19, 25%), and PNF (n=20, 26%) exercises, as well as 1 control group (n=19, 25%) with no intervention. Pressure pain threshold and the CPM responses at the myofascial trigger point, arm, and leg sites were assessed before and after the exercise session. The effective EIH response was reflected in the improvement of pressure pain thresholds.

Results: There was an increase in pressure pain thresholds and CPM responses at trigger point ($P<.001$ and $P<.001$), arm ($P<.001$ and $P<.001$), and leg sites ($P<.001$ and $P=.03$) in participants who performed PNF and isotonic exercise, while the isometric exercise only increased pressure pain thresholds at leg sites ($P=.03$). Compared with the control group, both the isotonic ($P=.02$) and PNF ($P<.001$) groups showed greater EIH responses at the trigger points. In comparison to the control group, only the PNF exercise ($P=.01$) significantly improved pressure pain thresholds and CPM responses at arm and leg sites compared to the control group.

Conclusions: PNF, isotonic, and isometric exercises could lead to local and global EIH effects. The improvement in CPM response following PNF and isotonic exercises suggested that the EIH mechanisms of different resistance exercises may be attributed to the enhancement of the endogenous pain modulation via the motor-sensory interaction from the additional eccentric and dynamic muscle contraction.

Trial Registration: Chinese Clinical Trial Registry ChiCtr202111090819166165; <https://tinyurl.com/2ab93p7n>

(*JMIRx Med* 2022;3(4):e40747) doi:[10.2196/40747](https://doi.org/10.2196/40747)

KEYWORDS

exercise induced hypoalgesia; proprioceptive neuromuscular facilitation; PNF; resistance exercise; conditioned pain modulation; myofascial pain syndrome; resistance training; hypoalgesia; exercise-induced hypoalgesia; shoulder myofascial pain; myofascial pain; pain management; chronic pain; musculoskeletal pain; physical therapy; physiotherapy; shoulder pain; upper back pain; exercise; pain

Introduction

Exercise therapy has been recommended for the management of chronic pain [1] regarding its advantages in efficiency, cost, safety, and the exercise-induced hypoalgesia effect (EIH) [2] in both healthy people and individuals with pain. Both aerobic and resistance exercises may attenuate pain perception globally by enhancing endogenous analgesia [3] via descending pain modulation. However, patients with myofascial pain syndrome (MPS) always demonstrate dysfunction in descending pain modulation [4], which could stimulate and disrupt pain perception and even lead to the attenuation of the hypoalgesia effect induced by maximal and submaximal aerobic [5] or resistance exercises [6].

The altered endogenous pain modulation may be the reason for impaired EIH for patients with MPS, because they showed both local and global hyperalgesia. Normally, the sufficient noxious or nonnoxious input from the C afferent fibers can activate the thalamic mediodorsal nucleus and ventromedial nucleus [7] at different thresholds, triggering the descending facilitation (low thresholds) or inhibition (high thresholds) [8], respectively. Moreover, the ON or OFF cells in the periaqueductal gray [9], locus coeruleus, and rostral ventromedial medulla [10], which are projected to the spinal dorsal horn, can either activate the N-methyl-D-aspartate receptors [11] to facilitate the ascending nociceptive signals or release the endogenous opioid [12], 5-hydroxytryptamine (5-HT) [13], and noradrenaline [14] to inhibit the nociceptive signals. Meanwhile, the pain-related brain regions, including the primary motor cortex [15], anterior cingulate cortex [16], and insula cortex [17], can also participate in the descending modulation of pain through the motor-sensory interaction. However, the prolonged pain condition, especially MPS, can develop into central sensitization, where the balance between the descending facilitation and inhibition is disrupted and may damage the EIH effects.

With respect to the pathway of endogenous pain modulation, it is believed that the sensory input during the exercise, especially

the proprioception and C fibers, can play a critical role in the EIH. Thus, proprioceptive neuromuscular facilitation (PNF) exercise [18], which can enhance the C fibers [19] and proprioception [20] inputs through the combination of various eccentric and dynamic exercises, may have additional positive effects on EIH in people with chronic pain; this is while the preliminary therapeutic effect of PNF on the pain of patients with MPS [21] has also been demonstrated. The type of muscle contraction can affect EIH in chronic pain. Specifically, isometric exercise could attenuate pain sensitivity in shoulder pain but not in fibromyalgia [22], while the isotonic exercise performed in nonpainful limbs could reduce pain perception in the case of chronic knee pain [23]. However, few studies have compared the effects of various resistance exercises on EIH in MPS. Moreover, the relationship between exercise type and endogenous analgesia features regarding EIH in MPS also needs further investigation.

In addition, conditioned pain modulation (CPM) [24] has been applied to evaluate the descending pain modulation in patients with chronic musculoskeletal pain. Additionally, the impairment of endogenous analgesia of MPS [25] was demonstrated by the attenuation of CPM responses. CPM has been shown to predict EIH in patients with knee osteoarthritis receiving bicycling and isometric exercise [26], while EIH has similar effects compared to CPM in healthy individuals [27]. However, the relationship between descending pain modulation and EIH has not yet been investigated in patients with MPS intervened by PNF and other resistance exercises.

Therefore, this pilot study aims to compare short-term EIH responses following PNF, isotonic, and isometric resistance exercises, as well as to investigate the relationship between EIH and descending pain modulation determined by CPM in patients with MPS. It was hypothesized that the PNF and all resistance exercises would have a higher EIH response compared with a blank control in individuals with MPS. The EIH and CPM response at baseline would be relatively lower in patients who have MPS with impaired descending pain modulation but would

increase or be restored following PNF and resistance exercise intervention performed on the affected areas.

Methods

Ethics Approval

This study has been approved by the Sports Science Experimental Ethics Committee of Beijing Sport University (ethics approval number: 2021153H) and registered in Chinese Clinical Trial Registry (registration number: ChiCtr202111090819166165).

Participants

A total of 76 female students (aged 18-30 years) from Beijing Sport University with shoulder MPS were enrolled in this study. This study selected female participants for the reason that EIH is potentially sex-related and is more consistently observed in women [28]. According to the pressure pain threshold (PPT) changes seen in previous studies [27,29], we used G-Power software with *F* test and ANCOVA parameters to calculate the sample size. The total samples of this study should be a minimum of 76 participants in the 3 groups, or 19 participants in each group.

The following inclusion criteria [30] were used to choose the participants: (1) reported shoulder pain persisting for at least 4 weeks up to 3 months and (2) had at least 1 latent trigger point on any side of the upper trapezius. The diagnosis of MPS adhered to the following standards: (1) palpation of a taut band, (2) identification of an exquisitely tender nodule (ie, the myofascial trigger points in the taut band), (3) reproduction of the patient's symptomatic pain with sustained pressure, and (4) the local twitch response. The threshold value of the visual analog scale (VAS) for the MPS is set at 30 mm/100mm. If multiple trigger points were detected in a participant, the trigger point with the lowest threshold of the pressure pain would be selected.

Individuals were excluded if they met the following standards: (1) confirmed or suspected spinal or shoulder injury, dislocation, and fracture or inflammatory or infective diseases; (2) had a history of spinal or shoulder surgery within 12 months, or other physical treatment within 1 month; and (3) presentation of cardiovascular conditions, psychosis, depression, cognitive impairment, or taking drugs for antidepressant or anticonvulsant

treatments, which would be carefully screened by a certified physician to ensure the safety of the intervention. The self-rating depression scale, self-rating anxiety scale, and brief psychiatric rating scale were applied and assessed by the physician during the screening periods.


All participants were randomly allocated into one of the four groups, as follows: group A (isometric exercise), group B (isotonic exercise), group C (PNF exercise), and group D (control). The randomized sequences were generated by a computer. All of the participants were labeled from number 01 to 76; then, the sequence was randomized using Excel software (Microsoft Corp) and allocated following the A-B-C-D circulation order. AN and XZH screened the participants.

Procedures

This study is designed as a randomized controlled trial. Participants who were included in this study were invited to perform exercise interventions of either isometric (group A), isotonic (group B), or PNF (group C) exercises, while the participants in control (group D) would rest for 15 minutes during the intervention session. Each exercise consists of 2 scapula movements and 1 shoulder movement. The intensity of exercise was set as 60% maximum voluntary contraction (MVC) to avoid pain in the context of significant analgesic effects seen in previous studies on EIH [27] and PNF [31-33], and it could be adjusted to the subpain threshold if participants reported pain during the exercise.

The CPM and EIH responses (measured by PPT of trigger point and other remote limbs) were assessed before and after the exercise session as outcome measurements. The VAS, height, weight and duration of shoulder pain were also collected before the intervention. The VAS was measured using a scale printed with a line ranging from 0 mm (no pain) to 100 mm (worst pain), and participants were asked to locate a point on the line to rate their current pain level. The VAS would be only measured as the baseline characteristics and would not be considered as an outcome measurement.

WZR and 2 other physical therapists evaluated all participants blinded to their exercise protocols. AN, XZH, and 1 other physical therapist guided all participants' exercise interventions. All of the participants received compensatory exercise and manual therapies following the outcome measurement and statistical analysis (Figure 1).

Figure 1. Flowchart of intervention. PNF: proprioceptive neuromuscular facilitation.

Interventions

Isometric Exercise

Participants in group A performed a modified isometric exercise program [34] including 2 scapula movements of scapula retraction (arm row at neutral position) and scapula elevation (dumbbell shrug at neutral position), and 1 shoulder abduction, where the dumbbells with adjustable weights were used. Scapula retraction and elevation were performed at 60% MVC or subpain threshold for 10 seconds with 15 sets at the neutral position. The shoulder isometric abduction was performed at 90°, and the elbow was flexed at 90°, with 60% MVC for 10 seconds of holding per 15 sets, and 30 seconds of rest was given between sets.

The MVCs of each movement were measured by a tension dynamometer with an LCD screen providing real-time statistics, where the participants were asked to perform a set of maximum voluntary contraction at the neutral position and adjust the contraction intensities by themselves during the exercise sessions. At 1 week before the intervention, this procedure was performed 3 times with a 30-second interval, and the mean value was identified as MVC.

Isotonic Exercise

Participants in group B performed a modified isotonic exercise program [35,36], including 2 scapula movements of dumbbell shrug and arm row and 1 shoulder lateral raise, where the dumbbells with adjustable weights were used. All of the isotonic exercises were performed at the moderate intensity [37] of 60% MVC or subpain threshold for 10 repetitions per 5 sets, where 1 minute of rest was given between sets.

PNF Exercise

Participants in group C performed a modified PNF exercise program [38,39], including the integration of agonist reversals (ARs), combination of isotonic contraction (CI), and rhythmic stabilization (RS) technique with the scapular pattern and upper extremity pattern. The upper extremity pattern D2 (flexion, abduction, and external rotation) was carried out using CI, where

the starting position of the shoulder was 180° internal rotation, 45° adduction, and 0° extension, and the ending position of the shoulder was 0° external rotation, 135° abduction, and 180° flexion. During the CI training, participants were asked to perform the concentric shoulder external rotation, abduction, and flexion to the ending position, and perform the eccentric movements back to the starting position.

The scapular pattern D2 (anterior descending and posterior evaluation) was carried out using AR followed by RS. During the AR training, the starting position of the scapular was maximum protraction and downward rotation, and the ending position of the scapular was maximum retraction and upward rotation. Participants were asked to perform the concentric scapula retraction and upward rotation to the ending position, then finish the concentric reversal movements back to the starting position. Additionally, the scapular was set at a neutral position during the RS training, when the participants were asked to maintain the stability of the scapular and confronting the resistances from the physical therapist. All of the PNF exercise sessions performed 10 repetitions per 5 sets at approximately 60% MVC or subpain threshold, with 1 minute of rest between sets.

Outcome Measures

PPTs of Trigger Points

PPTs of trigger points were measured by a quantitative sensory testing protocol [40] via a handheld pressure algometer (Baseline Dolorimeter, Fabrication Enterprises) with a 1 cm² metal probe and applied at a rate of 0.5 kg/s. PPT was measured in the trigger point located in the upper trapezius, which was labeled by a sterile marker. Participants were instructed to report as soon as they perceived a pain intensity by the VAS score of 40 out of 100 (Pain40) during pressure application; then, that threshold was recorded as PPT. This test was performed 1 minute before and after the intervention, while the difference of PPTs during the exercise session was recorded as the local EIH responses.

PPTs of Remote Sites

PPTs of remote sites were measured at the point (5 cm below the lateral condyle of humerus) of extensor carpus radialis (test point of arms) and the point (10 cm below the lateral femoral condyle) of peroneus longus (test point of legs) ipsilateral to the exercise limbs and were performed 1 minute before and after each exercise session. The difference of PPTs during the exercise session was recorded as the remote EIH responses.

Conditioned Pain Modulation

The CPM response was measured by quantitative sensory testing protocol [41], with the test stimulation applied by pressure, and conditioned stimulation applied by cold water immerse [42]. Participants first received pressure stimulation at the ipsilateral extensor carpus radialis and report a PPT at Pain40 as a test stimulus. Then, participants were instructed to immerse the contralateral hand into cold water at 8 °C for 1 minute and report the PPT at Pain40 when the pressure applied again at the threshold after 30 seconds of immersing. The difference between the 2 PPTs was recorded as the response of CPM.

Data Analysis

The main outcome of this study was the PPT of the trigger point, while the secondary outcomes were the PPT of the remote site and CPM responses. Normality of all data was assessed by means of the 1-sample Kolmogorov-Smirnov test. Difference in baseline data (height, weight, duration of pain, and VAS) between the groups was verified by the 1-way ANOVA test.

Table 1. Baseline characteristics (N=76)^a.

Measurements	A (n=18), mean (SD)	B (n=19), mean (SD)	C (n=20), mean (SD)	D (n=19), mean (SD)	P value
Age (years)	21.38 (2.50)	20.84 (1.92)	21.00 (1.81)	21.21 (2.41)	.95
Height (cm)	164.83 (3.75)	166.00 (4.53)	164.55 (4.21)	166.21 (4.44)	.61
Weight (kg)	55.03 (7.27)	57.32 (9.31)	54.75 (5.37)	59.57 (8.17)	.88
Duration of pain (week)	7.56 (2.83)	6.53 (2.95)	6.85 (2.52)	7.53 (3.01)	.54
VAS ^b (mm)	45.08 (12.80)	43.76 (15.89)	43.67 (13.78)	42.63 (8.06)	.18

^a1-way ANOVA; significant difference was set at $P < .05$.

^bVAS: visual analog scale.

The EIH and CPM Following Exercises

There was a significant increase in PPT at trigger point and arm site after isotonic ($P < .001$) and PNF exercises ($P < .001$), whereas the isometric exercise and the control group showed no difference compared to the baseline. The PPT at the arm sites significantly improved following PNF ($P < .001$) and isotonic

The 1-way ANCOVA was used to examine whether there was a significant difference within the 4 groups, considering PPT, EIH, and CPM at post exercise, while the pre-exercise measurements were set as covariates. The Bonferroni method was applied in the post hoc multiple comparison. The 2-tailed paired t test was used for the comparison within groups. All data were processed using SPSS, version 21.0 (IBM Corp), and the statistical significance was set at $P < .05$ for all tests.

Results

Baseline Characteristics

Of the 76 participants with MPS, 18 (24%) in group A completed the isometric exercise, 19 (25%) in group B completed the isotonic exercise, 20 (26%) in group C completed the PNF exercise, and 19 (25%) in group D finished a blank session. The average duration of shoulder MPS among participants was 7.56, 6.53, 6.85, and 7.53 weeks in group A, B, C, and D, respectively. The majority of participants ($n=48$, 63%) with MPS presented a duration greater than 6 weeks, and 33% ($n=25$) of them indicated over 8 weeks. Prior to the first exercise intervention, over half of the participants ($n=44$, 58%) had moderate-to-severe pain syndrome, with a VAS score higher than 40 mm/100mm. The baseline participant characteristics, including age ($P = .95$), height ($P = .61$), weight ($P = .88$), duration of pain ($P = .54$), and VAS ($P = .18$), did not present significant differences between the groups (Table 1).

exercises ($P < .001$), and the PPT at the leg sites also changed after isometric ($P = .03$), isotonic ($P = .03$), and PNF ($P < .001$) exercises.

A single session of isotonic ($P = .01$) and PNF ($P = .001$) exercises significantly improved the CPM responses, while the isometric exercise and the control group showed no difference compared to the baseline (Table 2 and Figure 2).

Table 2. Intervention results within groups (N=76)^a.

Outcome measure and group	Before exercise, mean (SD)	After exercise, mean (SD)	P value
PPT^b—trigger point (kg/cm²)			
A	2.72 (0.51)	2.80 (0.50)	.41
B	2.49 (0.35)	2.78 (0.28) ^c	<.001
C	2.38 (0.41)	3.19 (0.47) ^c	<.001
D	2.43 (0.35)	2.46 (0.37)	.42
PPT—arm (kg/cm²)			
A	2.64 (0.71)	2.72 (0.56) ^c	.13
B	2.42 (0.37)	2.68 (0.34) ^c	<.001
C	2.52 (0.44)	3.02 (0.47) ^c	<.001
D	2.41 (0.39)	2.49 (0.40)	.15
PPT—leg (kg/cm²)			
A	3.91 (0.70)	4.15 (0.76)	.03
B	3.81 (0.61)	4.12 (0.70) ^c	.03
C	3.74 (0.55)	4.63 (0.80) ^c	<.001
D	3.80 (0.50)	3.78 (0.49)	.78
CPM^d (kg/cm²)			
A	0.16 (0.30)	0.18 (0.26)	.72
B	0.17 (0.28)	0.33 (0.25) ^c	.01
C	0.14 (0.19)	0.38 (0.27) ^c	.001
D	0.18 (0.15)	0.18 (0.13)	.91


^aPaired *t* test; significant difference was set at $P < .05$.

^bPPT: pressure pain threshold.

^cSignificant changes.

^dCPM: conditioned pain modulation.

Figure 2. Results of pain pressure threshold (PPT), exercise-induced hypoalgesia, and conditioned pain modulation (CPM) following exercise interventions. PNF: proprioceptive neuromuscular facilitation.



The Effect of Exercise Type on EIH and CPM

For the PPT of the trigger point site, both the PNF ($P < .001$) and isotonic exercises ($P = .02$) showed significantly higher increase compared to control group. Meanwhile, PNF exercise had a greater effect than isotonic ($P < .001$) and isometric ($P < .001$) exercises. However, there was no significant difference between the isotonic and isometric exercises (Table 3).

For the PPT of remote sites, only the PNF exercises increased significantly higher thresholds compared to the control group ($P < .001$ and $P < .001$), isotonic exercises ($P = .01$ and $P = .004$), and the isometric exercises ($P < .001$ and $P = .002$) at both arm and leg sites. For the CPM, only PNF exercise attained a significantly greater responses compared to both control group ($P = .01$) and isometric exercise ($P = .02$; Table 3).

Table 3. Between-group comparison results of the intervention^a.

Outcome measures and intergroups	Mean difference	SE	P value	95% CI	
				Lower bound	Upper bound
PPT^b—trigger point (kg/cm²)					
A					
B	-0.144	0.093	.76	-0.397	0.109
C	-0.638 ^c	0.094	<.001	-0.894	-0.383
D	0.130	0.094	<.001	-0.126	0.385
B					
C	-0.495 ^c	0.090	<.001	-0.734	-0.243
D	0.274 ^c	0.090	.02	0.028	0.519
C					
D	0.768 ^c	0.089	<.001	0.526	1.010
PPT—arm (kg/cm²)					
A					
B	-0.123	0.086	.95	-0.351	-0.111
C	-0.389 ^c	0.085	<.001	-0.618	-0.160
D	0.060	0.086	>.99	-0.174	0.294
B					
C	-0.266 ^c	0.083	.01	-0.492	-0.040
D	0.183	0.084	.20	-0.045	0.411
C					
D	0.449 ^c	0.083	<.001	0.223	0.675
PPT—leg (kg/cm²)					
A					
B	-0.053	0.165	>.99	-0.500	0.395
C	-0.623 ^c	0.163	.002	-1.067	-0.180
D	0.285	0.165	.53	-0.162	0.733
B					
C	-0.570 ^c	0.160	.004	-1.006	-0.135
D	0.338	0.162	.25	-0.103	0.779
C					
D	0.908 ^c	0.160	<.001	0.473	1.344
CPM^d (kg/cm²)					
A					
B	-0.152	0.069	.19	-0.339	0.034
C	-0.212 ^c	0.068	.02	-0.397	-0.027
D	0.007	0.069	>.99	-0.181	0.194
B					
C	-0.061	0.067	>.99	-0.243	0.122
D	0.159	0.068	.14	-0.026	0.344

Outcome measures and intergroups	Mean difference	SE	P value	95% CI	
				Lower bound	Upper bound
C					
D	0.219 ^c	0.067	.01	0.036	0.402

^a1-way ANCOVA, adjusted by Bonferroni. Significant difference was set at $P < .05$.

^bPPT: pressure pain threshold.

^cSignificant changes.

^dCPM: conditioned pain modulation.

Discussion

Principal Findings

This pilot study investigated the local and remote responses of EIH and CPM after PNF, isotonic, and isometric resistance exercises for patients with MPS. Our findings mostly met what we previously hypothesized. PPT was increased at trigger point, arm, and leg sites when participants performed PNF and isotonic exercise, but was only increased at leg sites when participants performed isometric exercise. Compared with the control group, both isotonic and PNF groups showed significant greater EIH responses at the trigger points. However, only the PNF exercise significantly improved PPT at remote sites and CPM responds compared to the control group.

Changes in Pain Modulation

MPS, generally regarded as a typical chronic musculoskeletal pain [43], is mainly characterized by the presence of trigger points [44], which is a hypersensitive area that can be palpated in a muscle taut band. The trigger point is possibly induced by a continuous nociceptive stimulus from the local energy crisis of overused muscle fibers [45], followed by pain sensitization [46], which means the impairment of descending pain modulation. Thus, chronic MPS could attenuate both the CPM and the EIH effect. Vaegter et al [47] reported an altered effect of EIH in patients with chronic low back pain, where the acute 6-minute walk failed to induce EIH in patients with greater pain sensitivity. Chretien et al [5] also found that the deficits of EIH were related to the reduced CPM among adolescent girls with chronic pain.

This study found that both PNF and isotonic exercises significantly improved the CPM responses of MPS patients. It suggested that exercise with optimal intensity and type could affect central pain modulation and mediate neurotransmitters and cytokines. Exercises with various resistances may enhance the descending inhibition and reduce pain by activating the endocannabinoids, endogenous opioids, and 5-HT system. Crombie et al [48] reported that the serum endocannabinoids increased alongside the attenuation of pain sensitivity after resistance exercises in healthy individuals, while Bobinski et al [13] also found an increase in 5-HT in rostral ventromedial medulla through low-intensity exercise. Meanwhile, the activation of pain-related cortex regions following exercises may also mediate the pain processing of the thalamus and periaqueductal gray. Cumminford et al [49] found that the pain perception and facilitation of the thalamus was constrained by primary motor cortex stimulation, while Ellingson et al [50]

reported that the DLPFC function improved and was correlated with pain reduction in patients with fibromyalgia after cycling exercise. Interestingly, Lial et al [51] found that PNF exercise significantly improved DLPFC activation, which may indicate potential impacts on the central pain modulation, and that it still needs further investigation.

Changes in Pain Perception

In this study, EIH was demonstrated at the trigger point, arm, and leg sites for patients with MPS following PNF and isotonic exercises, which is consistent with prior research under various painful conditions. In most cases, resistance exercises with optimal intensities can induce global and local analgesia effects. First, the global analgesic effects induced by PNF and resistance exercises were also verified by Burrows et al [23]; these findings showed that the isotonic shoulder exercises of nonpainful limbs in patients with knee osteoarthritis effectively reduced pain, while Kuppens et al [52] confirmed EIH responses at leg sites after moderate-intensity shoulder extension exercises. Koltyn et al [53] also demonstrated the EIH responses elicited by contralateral isometric contraction in healthy individuals. Second, the isotonic and PNF exercises elicited greater EIH than isometric exercises, which is also found by Chung et al [54], showing that the isotonic exercises have superior EIH responses than isometric exercises in patients with chronic neck pain. Lastly, only isometric exercise failed to attenuate the pain perception of trigger point, which is also consistent with a previous systematic review by Bonello et al [55], indicating that there is no consistent evidence for EIH following isometric exercises in patients with chronic pain. Staud et al [56] also found that isometric exercise increased pain intensity in patients with fibromyalgia.

Neuromuscular exercises such as PNF have a significant therapeutic effect on many musculoskeletal pain conditions. According to a meta-analysis by Gao et al [18], PNF has more beneficial effects on pain relief and waist function improvement in patients with chronic lower back pain than other exercise interventions. Regarding chronic neck pain, Lytras et al [57] found that neuromuscular inhibition therapy combined with exercise intervention effectively reduced pain rating and improved neck function, while PNF also had a significant effect on knee osteoarthritis [58] and patellofemoral pain syndrome [59].

Potential Mechanisms of EIH

In this study, PNF exercise had a greater analgesic effect on MPS after intervention compared to control group and other exercises. This may be explained by the enhanced proprioception

and C fiber inputs from the additional eccentric [38] and dynamic muscle contractions. Although the activation of noxious C fibers following the overload exercise [19] may trigger the mechanical allodynia [60] and delayed onset muscle soreness, the sufficient nonnoxious C fiber input during the eccentric contraction with subpain threshold intensity in the PNF or even the isotonic exercise may still activate the descending inhibition via thalamus ventromedial nucleus. Stackhouse et al [61] compared the analgesic effect between the noxious electrical stimulation and the eccentric plantar flexor exercise with moderate intensity, and found that eccentric exercise induced both mechanical and thermal pain perception effectively. Apart from the enhanced proprioception and the C fiber inputs during the PNF exercise, the interaction between the participants and the researcher may also have positive effects on the proprioception inputs, considering the resistance provided by manual contact and the personal adaptation from the physical therapist. In such conditions, the intensity and direction of resistance can be adjusted more relevant to patients' perception.

Apart from the PNF exercise, only the isotonic exercise showed a greater change of PPT compared to control group at the trigger point, implying that the onset hypoalgesia effect from the CPM test may have contributed these changes during the test. The CPM test applied in this study, which provided cold stimulus, can also activate the C fiber afferent and the descending inhibition [41], which may have the overlap effect with the EIH responses. Thus, the relationship between the CPM test and the exercise requires further investigation.

Limitations

This study has several limitations. First, the indicators of the pain tests were limited; for instance, the PPT combined with thermal pain thresholds might better reflect the real neurophysiological aspects of musculoskeletal pain. Second, individuals with MPS differed in terms of pain duration and intensity, which may have an impact on the consistency of the results. Third, all of the participants were young female students, so the possible gender and aging difference of the pain processing should be considered in future studies. Lastly, the EIH effectiveness of moderate-intensity resistance exercise in this work was insufficiently investigated, and still needs to be evaluated comprehensively by increasing the variety of exercise types and duration length of the interventions.

Conclusions

In summary, PNF, isotonic, and isometric exercises could exert significant local and global EIH effects for patients with MPS, which may be influenced by the proprioception stimulus under the exercise types. The significant increases in CPM response after PNF and isotonic exercises indicated that the EIH mechanisms of these moderate-intensity exercises may involve the enhancement of the central descending inhibitory function. The findings of this study can serve as theoretical foundations for further studies focusing on central mechanisms of EIH, which could optimize the effect of exercise interventions for chronic pain in future clinical practices.

Acknowledgments

We would like to thank He-Jing Xu, Yang-Zhi Wu, Xi-Min Wu, Zi-Hang Gao, Wei Liu, Mao-Li Li, Li-Hui Qin, Zhen-Yan Chen, Cheng-Huan Hao, Bai-Hong Meng, and all the researchers who provided help and advice in our experiments.

Conflicts of Interest

None declared.

Multimedia Appendix 1

CONSORT-eHEALTH checklist (V 1.6.1).

[PDF File (Adobe PDF File), 342 KB - [xmed_v3i4e40747_app1.pdf](#)]

References

1. Sitges C, Velasco-Roldán O, Crespí J, García-Dopico N, Segur-Ferrer J, González-Roldán AM, et al. Acute Effects of a Brief Physical Exercise Intervention on Somatosensory Perception, Lumbar Strength, and Flexibility in Patients with Nonspecific Chronic Low-Back Pain. *J Pain Res* 2021;14:487-500 [FREE Full text] [doi: [10.2147/JPR.S274134](#)] [Medline: [33633462](#)]
2. Wewege MA, Jones MD. Exercise-Induced Hypoalgesia in Healthy Individuals and People With Chronic Musculoskeletal Pain: A Systematic Review and Meta-Analysis. *J Pain* 2021 Jan;22(1):21-31. [doi: [10.1016/j.jpain.2020.04.003](#)] [Medline: [32599154](#)]
3. Holmes SA, Kim A, Borsook D. The brain and behavioral correlates of motor-related analgesia (MRA). *Neurobiol Dis* 2021 Jan;148:105158 [FREE Full text] [doi: [10.1016/j.nbd.2020.105158](#)] [Medline: [33157210](#)]
4. Shah JP, Thaker N, Heimur J, Aredo JV, Sikdar S, Gerber L. Myofascial Trigger Points Then and Now: A Historical and Scientific Perspective. *PM R* 2015 Jul;7(7):746-761 [FREE Full text] [doi: [10.1016/j.pmrj.2015.01.024](#)] [Medline: [25724849](#)]
5. Chrétien R, Lavoie S, Chalaye P, de Vette E, Counil F, Dallaire F, et al. Reduced endogenous pain inhibition in adolescent girls with chronic pain. *Scand J Pain* 2018 Oct 25;18(4):711-717. [doi: [10.1515/sjpain-2018-0071](#)] [Medline: [30007060](#)]

6. Kuithan P, Heneghan NR, Rushton A, Sanderson A, Falla D. Lack of Exercise-Induced Hypoalgesia to Repetitive Back Movement in People with Chronic Low Back Pain. *Pain Pract* 2019 Sep;19(7):740-750. [doi: [10.1111/papr.12804](https://doi.org/10.1111/papr.12804)] [Medline: [31187932](https://pubmed.ncbi.nlm.nih.gov/31187932/)]
7. You H, Lei J, Niu N, Yang L, Fan X, Tjølsen A, et al. Specific thalamic nuclei function as novel 'nociceptive discriminators' in the endogenous control of nociception in rats. *Neuroscience* 2013 Mar 01;232:53-63. [doi: [10.1016/j.neuroscience.2012.12.021](https://doi.org/10.1016/j.neuroscience.2012.12.021)] [Medline: [23262232](https://pubmed.ncbi.nlm.nih.gov/23262232/)]
8. You H, Lei J, Pertovaara A. Thalamus: The 'promoter' of endogenous modulation of pain and potential therapeutic target in pathological pain. *Neurosci Biobehav Rev* 2022 Aug;139:104745. [doi: [10.1016/j.neubiorev.2022.104745](https://doi.org/10.1016/j.neubiorev.2022.104745)] [Medline: [35716873](https://pubmed.ncbi.nlm.nih.gov/35716873/)]
9. Lei J, Sun T, Lumb BM, You H. Roles of the periaqueductal gray in descending facilitatory and inhibitory controls of intramuscular hypertonic saline induced muscle nociception. *Exp Neurol* 2014 Jul;257:88-94. [doi: [10.1016/j.expneurol.2014.04.019](https://doi.org/10.1016/j.expneurol.2014.04.019)] [Medline: [24792920](https://pubmed.ncbi.nlm.nih.gov/24792920/)]
10. Sagalajev B, Viisanen H, Wei H, Pertovaara A. Descending antinociception induced by secondary somatosensory cortex stimulation in experimental neuropathy: role of the medullospinal serotonergic pathway. *J Neurophysiol* 2017 Mar 01;117(3):1200-1214 [FREE Full text] [doi: [10.1152/jn.00836.2016](https://doi.org/10.1152/jn.00836.2016)] [Medline: [28053243](https://pubmed.ncbi.nlm.nih.gov/28053243/)]
11. Da Silva LFS, Walder RY, Davidson BL, Wilson SP, Sluka KA. Changes in expression of NMDA-NR1 receptor subunits in the rostral ventromedial medulla modulate pain behaviors. *Pain* 2010 Oct;151(1):155-161 [FREE Full text] [doi: [10.1016/j.pain.2010.06.037](https://doi.org/10.1016/j.pain.2010.06.037)] [Medline: [20688433](https://pubmed.ncbi.nlm.nih.gov/20688433/)]
12. Kim Y, Byun J, Choi I. Effect of Exercise on μ -Opioid Receptor Expression in the Rostral Ventromedial Medulla in Neuropathic Pain Rat Model. *Ann Rehabil Med* 2015 Jun;39(3):331-339 [FREE Full text] [doi: [10.5535/arm.2015.39.3.331](https://doi.org/10.5535/arm.2015.39.3.331)] [Medline: [26161338](https://pubmed.ncbi.nlm.nih.gov/26161338/)]
13. Bobinski F, Ferreira TAA, Córdova MM, Dombrowski PA, da Cunha C, Santo CCDE, et al. Role of brainstem serotonin in analgesia produced by low-intensity exercise on neuropathic pain after sciatic nerve injury in mice. *Pain* 2015 Dec;156(12):2595-2606 [FREE Full text] [doi: [10.1097/j.pain.0000000000000372](https://doi.org/10.1097/j.pain.0000000000000372)] [Medline: [26447701](https://pubmed.ncbi.nlm.nih.gov/26447701/)]
14. Taylor BK, Westlund KN. The noradrenergic locus coeruleus as a chronic pain generator. *J Neurosci Res* 2017 Jun;95(6):1336-1346 [FREE Full text] [doi: [10.1002/jnr.23956](https://doi.org/10.1002/jnr.23956)] [Medline: [27685982](https://pubmed.ncbi.nlm.nih.gov/27685982/)]
15. Borovskis J, Cavaleri R, Blackstock F, Summers SJ. Transcranial Direct Current Stimulation Accelerates The Onset of Exercise-Induced Hypoalgesia: A Randomized Controlled Study. *J Pain* 2021 Mar;22(3):263-274. [doi: [10.1016/j.jpain.2020.08.004](https://doi.org/10.1016/j.jpain.2020.08.004)] [Medline: [32927091](https://pubmed.ncbi.nlm.nih.gov/32927091/)]
16. Zhou Y, Meng F, Cui Y, Xiong Y, Li X, Meng F, et al. Regular Aerobic Exercise Attenuates Pain and Anxiety in Mice by Restoring Serotonin-Modulated Synaptic Plasticity in the Anterior Cingulate Cortex. *Med Sci Sports Exerc* 2022 Apr 01;54(4):566-581 [FREE Full text] [doi: [10.1249/MSS.0000000000002841](https://doi.org/10.1249/MSS.0000000000002841)] [Medline: [34935710](https://pubmed.ncbi.nlm.nih.gov/34935710/)]
17. Zhang X, Zong B, Zhao W, Li L. Effects of Mind-Body Exercise on Brain Structure and Function: A Systematic Review on MRI Studies. *Brain Sci* 2021 Feb 07;11(2):205 [FREE Full text] [doi: [10.3390/brainsci11020205](https://doi.org/10.3390/brainsci11020205)] [Medline: [33562412](https://pubmed.ncbi.nlm.nih.gov/33562412/)]
18. Gao P, Tang F, Liu W, Mo Y. The effects of proprioceptive neuromuscular facilitation in treating chronic low back pain: A systematic review and meta-analysis. *J Back Musculoskelet Rehabil* 2022;35(1):21-33. [doi: [10.3233/BMR-200306](https://doi.org/10.3233/BMR-200306)] [Medline: [34250930](https://pubmed.ncbi.nlm.nih.gov/34250930/)]
19. Adreani CM, Hill JM, Kaufman MP. Responses of group III and IV muscle afferents to dynamic exercise. *J Appl Physiol* (1985) 1997 Jun;82(6):1811-1817 [FREE Full text] [doi: [10.1152/jappl.1997.82.6.1811](https://doi.org/10.1152/jappl.1997.82.6.1811)] [Medline: [9173945](https://pubmed.ncbi.nlm.nih.gov/9173945/)]
20. Areeudomwong P, Butttagat V. Proprioceptive neuromuscular facilitation training improves pain-related and balance outcomes in working-age patients with chronic low back pain: a randomized controlled trial. *Braz J Phys Ther* 2019;23(5):428-436 [FREE Full text] [doi: [10.1016/j.bjpt.2018.10.005](https://doi.org/10.1016/j.bjpt.2018.10.005)] [Medline: [30361077](https://pubmed.ncbi.nlm.nih.gov/30361077/)]
21. Sipko T, Glibowski E, Kuczyński M. Acute effects of proprioceptive neuromuscular facilitation exercises on the postural strategy in patients with chronic low back pain. *Complement Ther Clin Pract* 2021 Aug;44:101439. [doi: [10.1016/j.ctcp.2021.101439](https://doi.org/10.1016/j.ctcp.2021.101439)] [Medline: [34246129](https://pubmed.ncbi.nlm.nih.gov/34246129/)]
22. Lannersten L, Kosek E. Dysfunction of endogenous pain inhibition during exercise with painful muscles in patients with shoulder myalgia and fibromyalgia. *Pain* 2010 Oct;151(1):77-86. [doi: [10.1016/j.pain.2010.06.021](https://doi.org/10.1016/j.pain.2010.06.021)] [Medline: [20621420](https://pubmed.ncbi.nlm.nih.gov/20621420/)]
23. Burrows NJ, Booth J, Sturnieks DL, Barry BK. Acute resistance exercise and pressure pain sensitivity in knee osteoarthritis: a randomised crossover trial. *Osteoarthritis Cartilage* 2014 Mar;22(3):407-414 [FREE Full text] [doi: [10.1016/j.joca.2013.12.023](https://doi.org/10.1016/j.joca.2013.12.023)] [Medline: [24418672](https://pubmed.ncbi.nlm.nih.gov/24418672/)]
24. Ramaswamy S, Wodehouse T. Conditioned pain modulation-A comprehensive review. *Neurophysiol Clin* 2021 Jun;51(3):197-208. [doi: [10.1016/j.neucli.2020.11.002](https://doi.org/10.1016/j.neucli.2020.11.002)] [Medline: [33334645](https://pubmed.ncbi.nlm.nih.gov/33334645/)]
25. Vidor LP, Torres ILS, Medeiros LF, Dussán-Sarria JA, Dall'agnol L, Deitos A, et al. Association of anxiety with intracortical inhibition and descending pain modulation in chronic myofascial pain syndrome. *BMC Neurosci* 2014 Mar 19;15:42 [FREE Full text] [doi: [10.1186/1471-2202-15-42](https://doi.org/10.1186/1471-2202-15-42)] [Medline: [24645677](https://pubmed.ncbi.nlm.nih.gov/24645677/)]
26. Fingleton C, Smart KM, Doody CM. Exercise-induced Hypoalgesia in People With Knee Osteoarthritis With Normal and Abnormal Conditioned Pain Modulation. *Clin J Pain* 2017 May;33(5):395-404. [doi: [10.1097/AJP.0000000000000418](https://doi.org/10.1097/AJP.0000000000000418)] [Medline: [27518487](https://pubmed.ncbi.nlm.nih.gov/27518487/)]

27. Vaegter HB, Handberg G, Graven-Nielsen T. Similarities between exercise-induced hypoalgesia and conditioned pain modulation in humans. *Pain* 2014 Jan;155(1):158-167. [doi: [10.1016/j.pain.2013.09.023](https://doi.org/10.1016/j.pain.2013.09.023)] [Medline: [24076045](https://pubmed.ncbi.nlm.nih.gov/24076045/)]
28. Naugle KM, Naugle KE, Fillingim RB, Riley JL. Isometric exercise as a test of pain modulation: effects of experimental pain test, psychological variables, and sex. *Pain Med* 2014 Apr;15(4):692-701 [FREE Full text] [doi: [10.1111/pme.12312](https://doi.org/10.1111/pme.12312)] [Medline: [24308352](https://pubmed.ncbi.nlm.nih.gov/24308352/)]
29. Vaegter HB, Handberg G, Graven-Nielsen T. Isometric exercises reduce temporal summation of pressure pain in humans. *Eur J Pain* 2015 Aug;19(7):973-983. [doi: [10.1002/ejp.623](https://doi.org/10.1002/ejp.623)] [Medline: [25371064](https://pubmed.ncbi.nlm.nih.gov/25371064/)]
30. Fernández-de-Las-Peñas C, Dommerholt J. International Consensus on Diagnostic Criteria and Clinical Considerations of Myofascial Trigger Points: A Delphi Study. *Pain Med* 2018 Jan 01;19(1):142-150. [doi: [10.1093/pm/pnx207](https://doi.org/10.1093/pm/pnx207)] [Medline: [29025044](https://pubmed.ncbi.nlm.nih.gov/29025044/)]
31. Lim W. Optimal intensity of PNF stretching: maintaining the efficacy of stretching while ensuring its safety. *J Phys Ther Sci* 2018 Aug;30(8):1108-1111 [FREE Full text] [doi: [10.1589/jpts.30.1108](https://doi.org/10.1589/jpts.30.1108)] [Medline: [30154610](https://pubmed.ncbi.nlm.nih.gov/30154610/)]
32. Kwak DH, Ryu YU. Applying proprioceptive neuromuscular facilitation stretching: optimal contraction intensity to attain the maximum increase in range of motion in young males. *J Phys Ther Sci* 2015 Jul;27(7):2129-2132 [FREE Full text] [doi: [10.1589/jpts.27.2129](https://doi.org/10.1589/jpts.27.2129)] [Medline: [26310658](https://pubmed.ncbi.nlm.nih.gov/26310658/)]
33. Feland JB, Marin HN. Effect of submaximal contraction intensity in contract-relax proprioceptive neuromuscular facilitation stretching. *Br J Sports Med* 2004 Aug;38(4):E18 [FREE Full text] [doi: [10.1136/bjism.2003.010967](https://doi.org/10.1136/bjism.2003.010967)] [Medline: [15273211](https://pubmed.ncbi.nlm.nih.gov/15273211/)]
34. Price J, Rushton A, Tyros I, Tyros V, Heneghan NR. Effectiveness and optimal dosage of exercise training for chronic non-specific neck pain: A systematic review with a narrative synthesis. *PLoS One* 2020;15(6):e0234511 [FREE Full text] [doi: [10.1371/journal.pone.0234511](https://doi.org/10.1371/journal.pone.0234511)] [Medline: [32520970](https://pubmed.ncbi.nlm.nih.gov/32520970/)]
35. Dubé MO, Desmeules F, Lewis J, Roy J. Rotator cuff-related shoulder pain: does the type of exercise influence the outcomes? Protocol of a randomised controlled trial. *BMJ Open* 2020 Nov 05;10(11):e039976 [FREE Full text] [doi: [10.1136/bmjopen-2020-039976](https://doi.org/10.1136/bmjopen-2020-039976)] [Medline: [33154058](https://pubmed.ncbi.nlm.nih.gov/33154058/)]
36. Andersen LL, Kjaer M, Sjøgaard K, Hansen L, Kryger AI, Sjøgaard G. Effect of two contrasting types of physical exercise on chronic neck muscle pain. *Arthritis Rheum* 2008 Jan 15;59(1):84-91 [FREE Full text] [doi: [10.1002/art.23256](https://doi.org/10.1002/art.23256)] [Medline: [18163419](https://pubmed.ncbi.nlm.nih.gov/18163419/)]
37. da Cunha Ribeiro RP, Franco TC, Pinto AJ, Pontes Filho MAG, Domiciano DS, de Sá Pinto AL, et al. Prescribed Versus Preferred Intensity Resistance Exercise in Fibromyalgia Pain. *Front Physiol* 2018 Aug 10;9:1097 [FREE Full text] [doi: [10.3389/fphys.2018.01097](https://doi.org/10.3389/fphys.2018.01097)] [Medline: [30158876](https://pubmed.ncbi.nlm.nih.gov/30158876/)]
38. Oh D, Yoo K. The effects of therapeutic exercise using PNF on the size of calcium deposits, pain self-awareness, and shoulder joint function in a calcific tendinitis patient: a case study. *J Phys Ther Sci* 2017 Jan;29(1):163-167 [FREE Full text] [doi: [10.1589/jpts.29.163](https://doi.org/10.1589/jpts.29.163)] [Medline: [28210065](https://pubmed.ncbi.nlm.nih.gov/28210065/)]
39. Lee J, Park S, Na S. The effect of proprioceptive neuromuscular facilitation therapy on pain and function. *J Phys Ther Sci* 2013 Jun;25(6):713-716 [FREE Full text] [doi: [10.1589/jpts.25.713](https://doi.org/10.1589/jpts.25.713)] [Medline: [24259836](https://pubmed.ncbi.nlm.nih.gov/24259836/)]
40. Wyrażek M, Huber J, Lipiec J, Kulczyk A. Evaluation of palpation, pressure algometry, and electromyography for monitoring trigger points in young participants. *J Manipulative Physiol Ther* 2015;38(3):232-243. [doi: [10.1016/j.jmpt.2014.12.005](https://doi.org/10.1016/j.jmpt.2014.12.005)] [Medline: [25616692](https://pubmed.ncbi.nlm.nih.gov/25616692/)]
41. Mertens MG, Hermans L, Crombez G, Goudman L, Calders P, Van Oosterwijck J, et al. Comparison of five conditioned pain modulation paradigms and influencing personal factors in healthy adults. *Eur J Pain* 2021 Jan;25(1):243-256. [doi: [10.1002/ejp.1665](https://doi.org/10.1002/ejp.1665)] [Medline: [32965727](https://pubmed.ncbi.nlm.nih.gov/32965727/)]
42. Coulombe-Lévêque A, Tousignant-Laflamme Y, Léonard G, Marchand S. The effect of conditioning stimulus intensity on conditioned pain modulation (CPM) hypoalgesia. *Can J Pain* 2021 Feb 03;5(1):22-29 [FREE Full text] [doi: [10.1080/24740527.2020.1855972](https://doi.org/10.1080/24740527.2020.1855972)] [Medline: [33987521](https://pubmed.ncbi.nlm.nih.gov/33987521/)]
43. Saxena A, Chansoria M, Tomar G, Kumar A. Myofascial pain syndrome: an overview. *J Pain Palliat Care Pharmacother* 2015 Mar;29(1):16-21. [doi: [10.3109/15360288.2014.997853](https://doi.org/10.3109/15360288.2014.997853)] [Medline: [25558924](https://pubmed.ncbi.nlm.nih.gov/25558924/)]
44. Gerwin RD. Myofascial Trigger Point Pain Syndromes. *Semin Neurol* 2016 Oct;36(5):469-473. [doi: [10.1055/s-0036-1586262](https://doi.org/10.1055/s-0036-1586262)] [Medline: [27704503](https://pubmed.ncbi.nlm.nih.gov/27704503/)]
45. Simons DG. New views of myofascial trigger points: etiology and diagnosis. *Arch Phys Med Rehabil* 2008 Jan;89(1):157-159. [doi: [10.1016/j.apmr.2007.11.016](https://doi.org/10.1016/j.apmr.2007.11.016)] [Medline: [18164347](https://pubmed.ncbi.nlm.nih.gov/18164347/)]
46. Woolf CJ. Central sensitization: implications for the diagnosis and treatment of pain. *Pain* 2011 Mar;152(3 Suppl):S2-S15 [FREE Full text] [doi: [10.1016/j.pain.2010.09.030](https://doi.org/10.1016/j.pain.2010.09.030)] [Medline: [20961685](https://pubmed.ncbi.nlm.nih.gov/20961685/)]
47. Vaegter HB, Petersen KK, Sjødsholm LV, Schou P, Andersen MB, Graven-Nielsen T. Impaired exercise-induced hypoalgesia in individuals reporting an increase in low back pain during acute exercise. *Eur J Pain* 2021 May;25(5):1053-1063. [doi: [10.1002/ejp.1726](https://doi.org/10.1002/ejp.1726)] [Medline: [33400333](https://pubmed.ncbi.nlm.nih.gov/33400333/)]
48. Crombie KM, Brellenthin AG, Hillard CJ, Koltyn KF. Endocannabinoid and Opioid System Interactions in Exercise-Induced Hypoalgesia. *Pain Med* 2018 Jan 01;19(1):118-123 [FREE Full text] [doi: [10.1093/pm/pnx058](https://doi.org/10.1093/pm/pnx058)] [Medline: [28387833](https://pubmed.ncbi.nlm.nih.gov/28387833/)]
49. Cummiford CM, Nascimento TD, Foerster BR, Clauw DJ, Zubietta J, Harris RE, et al. Changes in resting state functional connectivity after repetitive transcranial direct current stimulation applied to motor cortex in fibromyalgia patients. *Arthritis Res Ther* 2016 Feb 03;18:40 [FREE Full text] [doi: [10.1186/s13075-016-0934-0](https://doi.org/10.1186/s13075-016-0934-0)] [Medline: [26842987](https://pubmed.ncbi.nlm.nih.gov/26842987/)]

50. Ellingson LD, Stegner AJ, Schwabacher IJ, Koltyn KF, Cook DB. Exercise Strengthens Central Nervous System Modulation of Pain in Fibromyalgia. *Brain Sci* 2016 Feb 26;6(1):8 [FREE Full text] [doi: [10.3390/brainsci6010008](https://doi.org/10.3390/brainsci6010008)] [Medline: [26927193](https://pubmed.ncbi.nlm.nih.gov/26927193/)]
51. Lial L, Moreira R, Correia L, Andrade A, Pereira AC, Lira R, et al. Proprioceptive neuromuscular facilitation increases alpha absolute power in the dorsolateral prefrontal cortex and superior parietal cortex. *Somatosens Mot Res* 2017 Sep;34(3):204-212. [doi: [10.1080/08990220.2017.1392298](https://doi.org/10.1080/08990220.2017.1392298)] [Medline: [29096587](https://pubmed.ncbi.nlm.nih.gov/29096587/)]
52. Kuppens K, Struyf F, Nijs J, Cras P, Franssen E, Hermans L, et al. Exercise- and Stress-Induced Hypoalgesia in Musicians with and without Shoulder Pain: A Randomized Controlled Crossover Study. *Pain Physician* 2016 Feb;19(2):59-68 [FREE Full text] [Medline: [26815250](https://pubmed.ncbi.nlm.nih.gov/26815250/)]
53. Koltyn KF, Umeda M. Contralateral attenuation of pain after short-duration submaximal isometric exercise. *J Pain* 2007 Nov;8(11):887-892. [doi: [10.1016/j.jpain.2007.06.003](https://doi.org/10.1016/j.jpain.2007.06.003)] [Medline: [17681886](https://pubmed.ncbi.nlm.nih.gov/17681886/)]
54. Chung S, Jeong Y. Effects of the craniocervical flexion and isometric neck exercise compared in patients with chronic neck pain: A randomized controlled trial. *Physiother Theory Pract* 2018 Dec;34(12):916-925. [doi: [10.1080/09593985.2018.1430876](https://doi.org/10.1080/09593985.2018.1430876)] [Medline: [29364754](https://pubmed.ncbi.nlm.nih.gov/29364754/)]
55. Bonello C, Girdwood M, De Souza K, Trinder NK, Lewis J, Lazarczuk SL, et al. Does isometric exercise result in exercise induced hypoalgesia in people with local musculoskeletal pain? A systematic review. *Phys Ther Sport* 2021 May;49:51-61. [doi: [10.1016/j.ptsp.2020.09.008](https://doi.org/10.1016/j.ptsp.2020.09.008)] [Medline: [33601254](https://pubmed.ncbi.nlm.nih.gov/33601254/)]
56. Staud R, Robinson ME, Price DD. Isometric exercise has opposite effects on central pain mechanisms in fibromyalgia patients compared to normal controls. *Pain* 2005 Nov;118(1-2):176-184. [doi: [10.1016/j.pain.2005.08.007](https://doi.org/10.1016/j.pain.2005.08.007)] [Medline: [16154700](https://pubmed.ncbi.nlm.nih.gov/16154700/)]
57. Lytras DE, Sykaras EI, Christoulas KI, Myrogiannis IS, Kellis E. Effects of Exercise and an Integrated Neuromuscular Inhibition Technique Program in the Management of Chronic Mechanical Neck Pain: A Randomized Controlled Trial. *J Manipulative Physiol Ther* 2020 Feb;43(2):100-113. [doi: [10.1016/j.jmpt.2019.03.011](https://doi.org/10.1016/j.jmpt.2019.03.011)] [Medline: [32482433](https://pubmed.ncbi.nlm.nih.gov/32482433/)]
58. Song Q, Shen P, Mao M, Sun W, Zhang C, Li L. Proprioceptive neuromuscular facilitation improves pain and descending mechanics among elderly with knee osteoarthritis. *Scand J Med Sci Sports* 2020 Sep;30(9):1655-1663. [doi: [10.1111/sms.13709](https://doi.org/10.1111/sms.13709)] [Medline: [32407583](https://pubmed.ncbi.nlm.nih.gov/32407583/)]
59. Motealleh A, Mohamadi M, Moghadam MB, Nejati N, Arjang N, Ebrahimi N. Effects of Core Neuromuscular Training on Pain, Balance, and Functional Performance in Women With Patellofemoral Pain Syndrome: A Clinical Trial. *J Chiropr Med* 2019 Mar;18(1):9-18 [FREE Full text] [doi: [10.1016/j.jcm.2018.07.006](https://doi.org/10.1016/j.jcm.2018.07.006)] [Medline: [31193229](https://pubmed.ncbi.nlm.nih.gov/31193229/)]
60. Nagi SS, Mahns DA. C-tactile fibers contribute to cutaneous allodynia after eccentric exercise. *J Pain* 2013 May;14(5):538-548. [doi: [10.1016/j.jpain.2013.01.009](https://doi.org/10.1016/j.jpain.2013.01.009)] [Medline: [23562300](https://pubmed.ncbi.nlm.nih.gov/23562300/)]
61. Stackhouse SK, Taylor CM, Eckenrode BJ, Stuck E, Davey H. Effects of Noxious Electrical Stimulation and Eccentric Exercise on Pain Sensitivity in Asymptomatic Individuals. *PM R* 2016 May;8(5):415-424. [doi: [10.1016/j.pmrj.2015.07.009](https://doi.org/10.1016/j.pmrj.2015.07.009)] [Medline: [26247163](https://pubmed.ncbi.nlm.nih.gov/26247163/)]

Abbreviations

- 5-HT:** 5-hydroxytryptamine
- AR:** agonist reversal
- CI:** combination of isotonic contraction
- CPM:** conditioned pain modulation
- EIH:** exercise-induced hypoalgesia
- MVC:** maximum voluntary contraction
- PNF:** proprioceptive neuromuscular facilitation
- PPT:** pressure pain threshold
- RS:** rhythmic stabilization
- VAS:** visual analog scale

Edited by E Meinert; submitted 04.07.22; peer-reviewed by J Greenberg, Anonymous, P Areudomwong; comments to author 17.11.22; revised version received 20.11.22; accepted 07.12.22; published 27.12.22.

Please cite as:

Xu ZH, An N, Wang ZR

Exercise-Induced Hypoalgesia Following Proprioceptive Neuromuscular Facilitation and Resistance Training Among Individuals With Shoulder Myofascial Pain: Randomized Controlled Trial

JMIRx Med 2022;3(4):e40747

URL: <https://med.jmirx.org/2022/4/e40747>

doi: [10.2196/40747](https://doi.org/10.2196/40747)

PMID: [37725522](https://pubmed.ncbi.nlm.nih.gov/37725522/)

©Zi-Han Xu, Nan An, Zi-Ru Wang. Originally published in JMIRx Med (<https://med.jmirx.org>), 27.12.2022. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in the JMIRx Med, is properly cited. The complete bibliographic information, a link to the original publication on <https://med.jmirx.org/>, as well as this copyright and license information must be included.

Publisher:
JMIR Publications
130 Queens Quay East.
Toronto, ON, M5A 3Y5
Phone: (+1) 416-583-2040
Email: support@jmir.org

<https://www.jmirpublications.com/>